

DISCUSSION PAPER SERIES

IZA DP No. 14695

**The Implications of Self-Reported Body
Weight and Height for Measurement
Error in BMI**

Apostolos Davillas
Andrew M Jones

AUGUST 2021

DISCUSSION PAPER SERIES

IZA DP No. 14695

The Implications of Self-Reported Body Weight and Height for Measurement Error in BMI

Apostolos Davillas

University of East Anglia and IZA

Andrew M Jones

University of York and Monash University

AUGUST 2021

Any opinions expressed in this paper are those of the author(s) and not those of IZA. Research published in this series may include views on policy, but IZA takes no institutional policy positions. The IZA research network is committed to the IZA Guiding Principles of Research Integrity.

The IZA Institute of Labor Economics is an independent economic research institute that conducts research in labor economics and offers evidence-based policy advice on labor market issues. Supported by the Deutsche Post Foundation, IZA runs the world's largest network of economists, whose research aims to provide answers to the global labor market challenges of our time. Our key objective is to build bridges between academic research, policymakers and society.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ISSN: 2365-9793

IZA – Institute of Labor Economics

Schaumburg-Lippe-Straße 5–9
53113 Bonn, Germany

Phone: +49-228-3894-0
Email: publications@iza.org

www.iza.org

ABSTRACT

The Implications of Self-Reported Body Weight and Height for Measurement Error in BMI*

We designed an experiment to explore the extent of measurement error in body mass index (BMI), when based on self-reported body weight and height. We find that there is a systematic age gradient in the reporting error in BMI, while there is limited evidence of systematic associations with gender, education and income. This is reassuring evidence for the use of self-reported BMI in studies that use it as an outcome, for example, to analyse socioeconomic gradients in obesity. However, our results suggest a complex structure of non-classical measurement error in BMI, depending on both individuals' and within-household peers' true BMI. This may bias studies that use BMI based on self-reported data as a regressor. Common methods to mitigate reporting error in BMI using predictions from corrective equations do not fully eliminate reporting heterogeneity associated with individual and within-household true BMI. Overall, the presence of non-classical error in BMI highlights the importance of collecting measured body weight and height data in large social science datasets.

JEL Classification: I10, C18, C50

Keywords: BMI, experiment, measurement error, reporting bias

Corresponding author:

Apostolos Davillas
Health Economics Group
Norwich Medical School
University of East Anglia
Norwich, NR4 7TJ
United Kingdom
E-mail: a.davillas@uea.ac.uk

* We are grateful to David Madden, Donal O'Neill, Nicolas Ziebarth, Robert Sugden and Stefan Penczynski for their comments on an earlier version of this paper. Understanding Society is an initiative funded by the Economic and Social Research Council and various Government Departments, with scientific leadership by the Institute for Social and Economic Research, University of Essex, and survey delivery by NatCen Social Research and Kantar Public. This paper makes use of data from the Understanding Society Innovation Panel administered by ISER, University of Essex, and funded by the Economic and Social Research Council. The research data are distributed by the UK Data Service. The funders, data creators and UK Data Service have no responsibility for the contents of this paper.

1 Introduction

Obesity is associated with increased risks of morbidity and mortality. This has led to a plethora of studies on the socio-economic consequences of obesity, such as labour market outcomes (Cawley, 2015). Because of the absence of measured anthropometric data in large-scale datasets, many existing studies are based on self-reports (Cawley, 2015; Cawley et al., 2015; Gil and Mora, 2011). The reliability of these measures in social science datasets is therefore of critical importance for obesity research.

We designed an experiment to explore the extent of measurement error in body mass index (BMI), when based on self-reported body weight and height, in the context of a multi-purpose survey. We collected information on self-reported body weight and height data immediately before the relevant physical measurements were taken.¹ The limited number of existing econometric analyses that examine measurement errors in anthropometrics mostly compare self-reports and measured anthropometric data that were collected with a considerable time difference and/or respondents were informed about the subsequent physical measurements (Cawley et al., 2015; Gil and Mora, 2011); these are also studies that are based on selected population samples (O’Neill and Sweetman, 2013).

The implications of measurement error are different depending on whether BMI is to be used as an outcome or as an explanatory variable. We explore whether the implied measurement error in BMI is systematically associated with socio-economic variables used in inequalities research. This is relevant for studies that use BMI as an outcome, modelled as a function of socioeconomic status (SES), and where measurement error contributes to the error term of the BMI regression equation. In addition, we explore whether the measurement error in BMI is non-classical, i.e., systematically associated with the measured values, and whether this association varies depending on the BMI of other household members. Non-classical measurement error may cause bias in regression models for other outcomes (e.g., earnings, health care costs) that use BMI as a regressor, even when instrumental variable methods are used to deal with endogeneity or errors-in-variables (O’Neill and Sweetman, 2013; Cawley et al., 2015).

As an extension, we revisit existing practices on using corrective equations² to partially address reporting error in weight and height in the absence of measured data (Cawley et al., 2015). We show that the predicted BMI values from these corrective equations still suffer from measurement error that depends on an individual’s own and within household peer’s measured BMI.

¹ The questionnaire design is structured so that respondents’ consent on the measurement followed their self-reports of weight/height and, thus, the latter is not contaminated by their informed consent to have their anthropometric measured.

² Corrective equations for self-reported body weight and height data rely on the external validity of the association between measured and reported weight/height values from one dataset to another. Corrections are based on the association in an external dataset that is translated to the main analysis dataset (“transferability” assumption).

2 Data

Understanding Society is a UK nationally representative household panel survey. One of its features is a sub-panel, the Innovation Panel (UKHLS-IP), reserved for experimental work.³

As part of the UKHLS-IP wave 12, we designed an experiment on the survey measurement of anthropometrics. Respondents were first asked for their self-reported body weight and height, followed by physical measurements⁴. The respondents gave their informed consent for these measurements (that follow conventional best practices on measurement of anthropometrics) at the point they were collected, which follows their self-reports of body weight/height. We focus on adults (aged 20+) here to eliminate any puberty-related body-size changes⁵.

Two BMI measures are calculated, as the body weight (Kg) divided by the square of height (metres), separately for the measured and the self-reported data. To facilitate interpretation of results, the absolute differences between the BMI based on self-reports and measured body weight and height data is used in our analysis (Cawley et al., 2015; Gil and Mora, 2011)⁶.

Our regression models for the absolute reporting error account for gender and age polynomials (in years divided by 10). Our SES measures are collected at UKHLS-IP wave 11: educational attainment (degree/post-secondary; A-level/equivalent; GCSE/equivalent; basic/no qualification) and household income (equivalised and log transformed). To explore the role of within-household peer effects, we use a dummy variable for being part of a household with low/moderate BMI levels (“low_hh_BMI”), defined as having an average BMI for all other adult household members, apart from the respondent, that is below the obesity threshold ($<30\text{kg/m}^2$)⁷.

³ The UKHLS-IP sample covers England, Wales and Scotland south of the Caledonian Canal.

⁴ Households were randomly allocated to two different survey modes to collect self-reports of body weight and height: a self-completion and an open interview mode. As we found no differences in reporting error by interview mode, these samples are pooled for our analysis.

⁵ To allow the results to be generalised to the population of Great Britain, we use sample weights that account for differential nonresponse, unequal selection during the sampling and non-response to our experiment. These weights are calculated by adjusting the UKHLS-IP wave 11 weights using a backward stepwise probit model on predictors from UKHLS-IP wave 12.

⁶ A limitation of raw reporting error is that under- and over-reports may cancel each other out and, thus, creating a misleading impression on the error’s magnitude. One may argue that under-reporting in BMI may be more important for a public health perspective as it may have more serious health consequences and result in an underestimation of the true overweight and obesity prevalence. However, the main scope of our analysis is to explore whether measurement error is non-classical. Given that the presence of non-classical error matters for models that use BMI as an explanatory variable (for example, wage equations, health care demand and costs), both under-reports and over-reports are of equal importance to get an unbiased estimate of the effect of adiposity on the outcome of interest.

⁷ Children are excluded here as (age and gender-specific) BMI in childhood should be interpreted differently than adult obesity. Our results are robust to a sensitivity analysis restricting our

3 Methods

Absolute reporting error is modelled by linear regression. Regression models are first estimated using the set of demographics and SES. To explore whether measurement error is non-classical, we add an individual’s own BMI based on their measured data. This specification is augmented by adding BMI information for the other household members and its interaction with an individual’s own (measured) BMI⁸.

As an extension, we test whether the conventional method of using corrective equations for self-reports of body weight/height is sufficient to mitigate reporting error and, more, importantly its systematic association with covariates (Cawley, 2015).⁹ Availability of self-reported and measured data allow us to estimate analogous corrective equations by regressing measured weight and height data on self-reports and a vector of demographics. To mimic correction procedures for self-reported anthropometrics in the existing studies, the predictions from these equations are used to calculate self-reports of body weight and height that are corrected for reporting error. To explore the remaining reporting error following this correction procedure we compute the absolute difference between the corrected and measured BMI. This measure of the remaining reporting error is regressed on our set of demographics, SES and individual’s own and within-household peer’s (measured) BMI to explore whether there are still systematic associations with these factors.

4 Results


Figure 1 shows that there is a high correlation between reported and measured BMI data. However, there is not a perfect match – reporting error is more likely to result in under-reporting of BMI than over-reporting; more of the data points are concentrated above, rather than below, the 45-degree equality line. Despite the small differences that initial visualisation in Figure 1 shows, obesity prevalence is systematically higher when based on measured (36% and 32% for females and males) as opposed to self-reported data (32% and 26% for females and males); this is evident in Table A1 (Appendix).

sample to those households with two and more household members (results available upon request). BMI values above the obesity threshold may be of particular interest here as they are more visible in people’s silhouettes and, thus, more likely to exert peer-effects on reporting behaviour (Lønnebotn et al., 2018).

⁹ In the existing economic studies of obesity that rely on self-reports only, corrective equations – based on the relationship between measured and self-reported body weight and height data – are estimated using alternative, rather than the main analysis, data source. Then, the coefficients from these equations are transferred to the analysis sample and, after multiplying the coefficients by the self-report values, they obtain measures of weight and height corrected for the reporting error (Cawley, 2015).

To further quantify the magnitude of reporting error, the mean of raw reporting errors (defined as reported minus measured BMI) show that, on average, respondents over-report their height (by 1.196 cm), under-report their weight (by 0.941 kg) and, consequently, BMI is underestimated by 0.738kg/m^2 (Tables 1 and Table A.2, Appendix). Our absolute measure of the reporting error shows that both under- and over-reports result in an average total error in BMI of 1.3kg/m^2 , i.e., 4.4% of measured BMI (Table 1). Graphs of the distributions of reporting error in weight, height and BMI are presented in Figure A.1 (Appendix).

Figure 1. Scatter plots of measured versus reported BMI.



Notes: Markers are scaled to reflect sample weights. Darker regions representing more concentrated data points. The black line is a 45-degree line.

Table 1. Summary statistics for measurement error in BMI(kg/m^2).

	Mean	Std. Dev.
Raw error	-0.738	1.541
Absolute error	1.266	1.147
Absolute error (%measured)	4.418	3.927

4.1. Regression analysis

Table 2 presents regression analyses of the absolute BMI reporting error. We find a non-linear and systematic association between age and the absolute reporting error in BMI across all model specifications, which increases steeply for those aged 70 and above (Figure 2). No systematic associations are evident for gender and SES. If BMI is the outcome of interest, for example, in an analysis of socioeconomic inequality, then we do not find systematic reporting error by SES.

Specification 2 shows that measurement error in BMI is non-classical, with the respondent’s own measured BMI being positively associated with the absolute error in BMI. Conditional on the individuals’ own BMI, their within-household peers BMI also plays an important role (specification 3). Specifically, as illustrated in Figure 3, although the predicted absolute error in self-reported BMI increases in magnitude for every unit increase in an individual’s measured BMI *ceteris paribus*, there is heterogeneity related to household-peer effects as suggested by the interaction term (Table 2). Respondents with measured BMI of around 31 and above, which coincides with the obesity threshold, reported anthropometrics more accurately (lower reporting error in BMI) when living in households with other members having low or moderate BMI values as opposed to those living in households with excess BMI levels.¹⁰

Sensitivity analysis shows that the time of anthropometric measurement during the day does not affect our results in Table 2; main effect of the time of the day and its interaction terms with measured BMI are not statistically significant (p -values >0.10).¹¹

4.2. Correction equations

Table 3 presents regression analysis on the absolute difference between BMI based on predictions from the corrective equations (Table A.3, Appendix) and measured BMI. Measured BMI still plays a systematic role in the remaining error in BMI (after the correction). Moreover, we observe similar patterns (but with lower magnitude of predicted errors) for the heterogeneous role of individuals’ measured BMI (on the remaining reporting error in BMI) related to household-peer effects to those observed in Figure 3, without adjustments using the corrective equations.

¹⁰ Overall, this suggests the presence of another source of non-classical measurement error, known as differential measurement error (O’Neill and Sweetman, 2013). Typically, differential measurement error is defined as another type of non-classical measurement error that arises when the reported measurement error in BMI is correlated with the error term in econometric models for which BMI is the independent variable of interest (O’Neill and Sweetman, 2013). For example, assume that the within-household peers’ BMI matters for the reporting error in individual’s BMI, as we will explore here. If the within-household peers’ BMI is directly or indirectly correlated with the outcome of interest (often, in our case, labour market outcomes, healthcare demand etc.) and, thus, captured by the regressions’ residuals, then, differential reporting error exists.

¹¹ Additional sensitivity analysis shows that our results on the role of within-household peer effects on the absolute measurement error in BMI are robust to conditioning for the number of household members responded and provided body weight and height data.

Table 2. Regression analysis of absolute BMI reporting error.

	Specification 1	Specification 2	Specification 3
Age	1.040 (0.651)	0.860 (0.703)	1.067 (0.686)
Age ²	-0.247** (0.126)	-0.216 (0.133)	-0.257** (0.130)
Age ³	0.018** (0.008)	0.017** (0.008)	0.019** (0.008)
Male	-0.019 (0.090)	0.007 (0.086)	0.006 (0.085)
Degree/post-secondary	-0.277* (0.156)	-0.208 (0.156)	-0.232 (0.156)
A-level/equivalent	-0.023 (0.185)	0.018 (0.182)	0.011 (0.180)
GCSE/equivalent	-0.109 (0.165)	-0.161 (0.160)	-0.170 (0.157)
Income	-0.003 (0.076)	-0.024 (0.074)	-0.021 (0.074)
BMI measured		0.049*** (0.010)	0.066*** (0.011)
Low_hh_BMI			1.586** (0.621)
BMI measured*Low_hh_BMI			-0.051** (0.022)
R-squared	0.049	0.117	0.137

*p<0.10;**p<0.05;***p<0.001.

Figure 2. Prediction (based on specification 3, Table 2) of the absolute BMI error by age.




Figure 3. Prediction (based on specification 3, Table 2) of the absolute error in self-reported BMI by measured BMI values and household BMI levels.





Table 3. Regression analysis of the absolute remaining BMI measurement error following adjustments from corrective equations.

	Coeff. (std. error)
Age	0.285 (0.615)
Age ²	-0.061 (0.119)
Age ³	0.004 (0.007)
Male	0.026 (0.078)
Degree/post-secondary	-0.098 (0.140)
A-level/equivalent	0.105 (0.156)
GCSE/equivalent	-0.054 (0.138)
Income	-0.106 (0.070)
BMI measured	0.043*** (0.010)
Low_hh_BMI	1.157** (0.584)
BMI measured*Low_hh_BMI	-0.036* (0.020)
R-squared	0.061

*p<0.10;**p<0.05;***p<0.001.

Figure 4. Prediction (based on Table 3) of the remaining BMI error by measured BMI values and household BMI levels.



5 Conclusion

We designed an experiment to measure reporting error in BMI. We find a systematic age gradient in the reporting error in BMI, while there is limited evidence of systematic associations with gender and SES. This is reassuring evidence for the use of self-reported BMI in studies that use it as an outcome, for example, to analyse socioeconomic gradients in obesity.

Reporting error in BMI is associated with individual's measured BMI. The role of an individual's measured BMI on reporting error varies as a result of within-household peer-effects: for individual's with measured BMI values above the obesity threshold, measurement error is higher for those living in households with other members having high BMI levels. The latter is broadly consistent with the role of social norms on health reporting (Gil and Mora, 2011) and challenges the between-households (or and above between-individuals) reliability of the self-reported data. This complex structure of non-classical measurement error may be an issue in studies that use self-reported BMI as an explanatory variable.

Common methods to mitigate reporting error in BMI using corrective equations not to fully eliminate systematic associations with individual and within-household BMI. Overall, as the error in anthropometrics is non-classical our results highlight the importance of collecting measured body weight and height data in large social science datasets.

References

- Cawley, J., Maclean, J.C., Hammer, M., Wintfeld, N.(2015). Reporting error in weight and its implications for bias in economic models. *Economics & Human Biology*, 19, 27-44.
- Cawley, J.(2015). An economy of scales: a selective review of obesity's economic causes, consequences, and solutions. *Journal of Health Economics*, 43, 244-268.
- Gil, J., Mora, T.(2011). The determinants of misreporting weight and height: the role of social norms. *Economics & Human Biology*, 9, 78-91.
- Lønnebotn, M., Svanes, C., Iglund, J., Franklin, K. A.,... & Demoly, P. (2018). Body silhouettes as a tool to reflect obesity in the past. *PloS ONE*, e0195697.
- O'Neill, D., Sweetman, O.(2013). The consequences of measurement error when estimating the impact of obesity on income. *IZA Journal of Labor Economics*, 2, 1-20.

Appendix

Table A.1 Classification of obesity using self-reported and measured BMI.

	Females	Males
Obesity prevalence (measured BMI)	36.43	32.16
Obesity prevalence (reported BMI)	32.42	26.10
P-value (difference)	0.003	0.000
Percentage classified as:		
True positive	30.51	24.37
False positive	1.92	1.73
True negative	61.65	66.11
False negative	5.93	7.79
Total	100.0	100.0
Sensitivity	83.7	75.8
Specificity	97.0	97.4

Table A.2 Reporting error in body weight and height.

	Mean	Std. Dev.
Body height (in cm)		
Raw error	1.196	2.767
Absolute error	2.243	2.013
Absolute error (% of measured)	1.344	1.228
Body weight (in kg)		
Raw error	-0.941	3.440
Absolute error	2.310	2.715
Absolute error (% of measured)	2.889	3.380
Sample size	873	

Table A.3 Prediction equation for measured body weight and height.


	Body weight (measured)	Body height (measured)
Body weight reported	1.008*** (0.012)	—
Body height reported	—	0.889*** (0.017)
Age†	-2.590 (2.691)	0.983*** (0.333)
Age ² †	0.371 (0.509)	-0.143*** (0.031)
Age ³ †	-0.015 (0.030)	—
Male	-18.293** (7.176)	0.898*** (0.323)
Male*Age	10.102** (4.449)	—
Male* Age ²	-1.772** (0.872)	—
Male*Age ³	0.098* (0.054)	—
Constant	5.814 (4.463)	16.453*** (3.018)
R-squared	0.969	0.937

Notes: Robust standard errors in parentheses.

†Age is divided by 10.

*p<0.10; **p<0.05; ***p<0.001.

Figure A1. Raw reporting error in weight, height, and BMI.



Notes: The red lines show the distribution of the raw reporting error in height, weight and BMI. Superimposed in each histogram is the corresponding normal distribution (black lines). In each graph, the vertical axis is the percentage of the sample and the horizontal axis represents units of raw measurement error.